

DISCLAIMER. This collection of practice problems is *not* guaranteed to be identical, in length or content, to the actual exam. You may expect to see problems on the test that are not exactly like problems you have seen before.

On the actual exam you will have more room to work the problems. You will see directions similar to these:

1. Please read directions carefully. Raise your hand if you are not sure what a problem is asking.
2. *You must explain your work thoroughly and unambiguously to receive full credit on questions or parts of questions designated as **Work and Answer**.*
3. **No calculators or notes are allowed on this exam.**
4. You have 2 hours to complete your test, unless announced otherwise. Do not spend too long on any one problem. You do not have to do the problems in order. Do the easy ones first. Do not attempt the bonus question until you have completed the rest of the test. Before turning in your test, please make sure you have answered and double-checked all the questions.
5. If you need scratch paper, please raise your hand. You may not use your own paper. When you have finished your exam, please turn in any scratch paper you use.
6. For **Work and Answer** problems, write your solutions in the space provided for each problem, or provide specific instructions as to where your work is to be found. *Make it clear what you want and don't want graded.* Your final answers should be boxed or circled.
7. Don't stress! I'm rooting for you!

You will also see the following useful formulas:

English system formulas:

$$1 \text{ ft.} = 12 \text{ in.}$$

$$5280 \text{ ft.} = 1 \text{ mi.}$$

$$16 \text{ oz.} = 1 \text{ lb.}$$

$$\int \sin^2 x \, dx = \frac{1}{2}x - \frac{1}{4}\sin 2x + C$$

$$\int \frac{1}{a^2 + x^2} \, dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + C$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} \, dx = \sin^{-1} \left(\frac{x}{a} \right) + C$$

$$\int a^x \, dx = \frac{1}{\ln a} a^x + C$$

$$\int \frac{\sqrt{a^2 + x^2}}{x} \, dx = \sqrt{a^2 + x^2} - a \ln \left| \frac{a + \sqrt{a^2 + x^2}}{x} \right| + C$$

(Formulas for volume, arc length, surface area, hydrostatic force, and center of mass)

Metric system formulas:

$$F = m \cdot a$$

$$g = 9.8 \text{ m/s}^2$$

$$100 \text{ cm} = 1 \text{ m}$$

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$\sin 2\theta = 2 \sin \theta \cos \theta$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta$$

Multiple Choice. Circle the letter of the best answer.

1. The area between the curves $f(x) = 3x - 1$ and $g(x) = x^2 + 1$ from $x = 2$ to $x = 5$ is

(a) $\int_2^5 (3x - 1) - (x^2 + 1) \, dx$

(c) $\int_2^3 (3x - 1) - (x^2 + 1) \, dx + \int_3^5 (x^2 + 1) - (3x - 1) \, dx$

(b) $\int_2^5 (x^2 + 1) - (3x - 1) \, dx$

(d) $\int_2^4 (x^2 + 1) - (3x - 1) \, dx + \int_4^5 (3x - 1) - (x^2 + 1) \, dx$

2. The weight of a leaky bucket when carried x feet up a 20-foot ladder is $30 - 2x$ pounds. The work done in carrying the bucket from the ground to the top of the ladder is

(a) 200 ft.-lb. (c) 300 ft.-lb.

(b) 250 ft.-lb. (d) 350 ft.-lb.

3. The average value of the function $f(x) = \sqrt[3]{x}$ from $x = 1$ to $x = 8$ is

(a) $\frac{45}{4}$ (c) $\frac{45}{28}$

(b) $\frac{45}{7}$ (d) $\frac{45}{14}$

4. $\int_0^1 x e^{3x} dx =$

(a) $\frac{1}{6}e^3$ (c) $\frac{1}{3}(e^3 - 1)$

(b) $\frac{2}{3}(e^3 + 1)$ (d) $\frac{1}{9}(2e^3 + 1)$

5. $\int \sin^3 x \cos x dx =$

(a) $\frac{1}{4} \cos^4 x + C$ (c) $-\frac{1}{4} \sin^4 x + C$

(b) $\frac{1}{4} \sin^4 x + C$ (d) $-\frac{1}{4} \cos^4 x + C$

6. Using trigonometric substitution, the integral $\int \frac{x^3}{\sqrt{1-x^2}} dx$ is equal to

(a) $\int \sin^3 \theta d\theta$ (c) $\int \frac{\sin^3 \theta}{\cos^2 \theta} d\theta$

(b) $\int \frac{\sin^3 \theta}{\cos \theta} d\theta$ (d) $\int \sin^2 \theta \cos \theta d\theta$

7. $\int \frac{3}{2+x^2} dx =$

(a) $\frac{3}{2} \tan^{-1}(x) + C$ (c) $\frac{3\sqrt{2}}{2} \tan^{-1}\left(\frac{x}{\sqrt{2}}\right) + C$

(b) $-\frac{3}{2+x} + C$ (d) $-\frac{3}{4}x^2 + C$

8. $\int_2^\infty \frac{1}{x^4} dx =$

(a) $\frac{1}{6}$

(c) $\frac{3}{20}$

(b) $\frac{1}{24}$

(d) ∞ (diverges)

9. The length of the curve $f(x) = 5x^2$ from $x = 1$ to $x = 4$ is

(a) $\int_1^4 \sqrt{1 + 100x^2} dx$

(c) $\int_1^4 \sqrt{1 + 5x^2} dx$

(b) $\int_1^4 1 + 5x^2 dx$

(d) $\int_1^4 \sqrt{10x} dx$

10. The area of the surface formed by rotating the curve $f(x) = 5x^2$ from $x = 1$ to $x = 4$ about the y -axis is

(a) $\int_1^4 \sqrt{1 + 100x^2} dx$

(c) $2\pi \int_1^4 \sqrt{1 + 5x^2} dx$

(b) $2\pi \int_1^4 x\sqrt{1 + 100x^2} dx$

(d) $\int_1^4 5x^2\sqrt{10x} dx$

11. After eliminating the parameter, the curve $\begin{matrix} x = e^t - t \\ y = t^3 \end{matrix}$ is identical to the curve

(a) $x = e^{\sqrt[3]{y}} - \sqrt[3]{y}$

(c) $y = (e^x - x)^3$

(b) $x = e^{y^3} - y^3$

(d) $y = e^{x^3} - x^3$

12. The Cartesian coordinates for the polar point $(-\frac{1}{2}, \frac{3\pi}{2})$ are

(a) $(\frac{1}{2}, 0)$

(c) $(-\frac{1}{2}, 0)$

(b) $(0, \frac{1}{2})$

(d) $(0, -\frac{1}{2})$

13. The sequence $a_n = \frac{2n}{3n-1}$

(a) converges to 0

(c) converges to $\frac{2}{3}$

(b) converges to 1

(d) diverges

14. The series $\sum_{n=1}^{\infty} \frac{2n}{3n-1}$

(a) converges to 0

(c) converges to $\frac{2}{3}$

(b) converges to 1

(d) diverges

15. In order to determine whether or not the series $\sum_{n=1}^{\infty} \frac{(-1)^n 5n}{3n^2 - 1}$ converges, the limit comparison test may be used with comparison series $\sum b_n =$

(a) $\sum \frac{5}{3n^2}$

(c) $\sum \frac{1}{n^2}$

(b) $\sum \frac{(-1)^n}{n}$

(d) none; the limit comparison test cannot be used

16. The series $\sum_{n=0}^{\infty} \frac{(-1)^n (x-2)^n}{5^n}$

(a) converges for $\frac{9}{5} < x < \frac{11}{5}$ only (c) converges for all x

(b) converges for $-3 < x < 7$ only (d) converges only for $x = 2$ only

17. The series $\sum_{n=0}^{\infty} \frac{(-1)^n 3^{2n}}{(2n)!}$

(a) converges to $\sin 3$

(c) converges to 3

(b) converges to $\cos 3$

(d) diverges

18. The binomial series of the function $f(x) = \frac{1}{(1-x)^5}$ is

(a) $\sum_{n=0}^{\infty} \binom{-5}{n} (-1)^n x^n$

(c) $\sum_{n=0}^{\infty} \binom{5}{n} (-x)^n$

(b) $\sum_{n=0}^{\infty} \binom{-5}{n} x^n$

(d) $\sum_{n=0}^{\infty} \binom{n}{-5} (-1)^n x^n$

Fill-In.

1. *Circle the best answer.*

In order to find the volume of the solid formed by rotating the region enclosed by $y = x^2 + 1$ and $y = 3x - 1$ about the x -axis, it is best to use the (**disk** | **shell**) method.

2. *Fill in the correct numerators. Your answers should be numbers or polynomials, not A, B, etc.*

$$\frac{4x - 1}{(x - 2)(x + 3)^2} = \frac{\boxed{}}{x - 2} + \frac{\boxed{}}{x + 3} + \frac{\boxed{}}{(x + 3)^2}$$


3. Circle the best answer.

The series $\sum_{n=1}^{\infty} \frac{3n^2 - 1}{n^5}$ (converges absolutely | converges conditionally | diverges)

4. $\binom{3/2}{4} = \frac{\quad}{(number)}$.

Graphs. More accuracy = more points!


1. On the axes at right, sketch the solid formed by rotating the region shown about the y -axis.



(a) For the function $f(x)$ graphed at right, sketch a **rectangle** on the same axes whose area is approximately $\int_1^5 f(x) dx$.

2. (b) The average value f_{ave} of $f(x)$ from $x = 1$ to $x = 5$ is approximately _____ .

(c) The approximate value(s) of c so that $f(c) = f_{\text{ave}}$ is/are _____ (list all values).



Work and Answer. You must show all relevant work to receive full credit.

1. Find the area of the region enclosed by the curves $y = 4 - x^2$ and $y = x + 2$.

2. If the force required to pump water at depth x over the side of a tank 2 meters deep is $F(x) = 9800(3x^2 + 4x)$ Newtons, find the work done to pump all the water out.

3. Evaluate $\int \ln x dx$.

4. Evaluate $\int \tan^{-1} x \, dx$.

5. Evaluate $\int_0^{\infty} x e^{-x^2} \, dx$.

6. Find the sum $\sum_{n=0}^{\infty} \frac{3}{4^n}$.

7. Determine whether the series $\sum_{n=0}^{\infty} \frac{(-1)^n (n^2 + 1)}{5^n}$ is absolutely convergent, conditionally convergent, or divergent.

8. Find the interval of convergence of the power series $\sum_{n=1}^{\infty} \sqrt{n} x^n$.

9. Find a power series representation for the function $f(x) = \ln(1 + x)$.

10. Find a power series representation for the function $f(x) = \sin(2x)$.

11. (a) Find a power series representation for the function $f(x) = (2 + 3x)^{10}$.
(b) Find the coefficient of x^3 in the above series.

Some kind of **BONUS**.