

### Finding the $n$ -th term of a sequence - Section 11.1

To develop a formula for the  $n$ -th term of a sequence  $\{a_n\}$ , look for the following patterns:

(a) **The Arithmetic Pattern.**

If the terms count up by the same number each time, then the  $n$ -th term will look like that number times  $n$ , plus or minus some correction term.

**Example.**  $\{-4, -1, 2, 5, 8, 11, \dots\}$ .

Notice that each term in the sequence is 3 more than the one before. Therefore the  $n$ -th term is  $3n \pm \xi$ , where the  $\xi$  depends on where the sequence starts. If the first term is  $a_1$ , then the  $n$ -th term is  $a_n = 3n - 7$ , since  $a_1$  must be  $-4$ . If the first term is  $a_0$ , then the  $n$ -th term is  $a_n = 3n - 4$ , since  $a_0$  must be  $-4$ .

The following are all correct representations of the above sequence:

- $\{3n - 7\}_{n=1}^{\infty}$
- $\{3n - 4\}_{n=0}^{\infty}$
- $\{3n + 2\}_{n=-2}^{\infty}$

(b) **The Power Pattern.**

If the terms are obtained by taking powers of successive integers, then the  $n$ -th term will look like  $(n \pm \xi)^{\bullet}$ , where the  $\xi$  depends on where the sequence starts.

**Example.**  $\{4, 9, 16, 25, \dots\}$ .

Notice that each term in the sequence is a perfect square, starting with  $2^2$ . Therefore the  $n$ -th term is  $(n \pm \xi)^2$  for some  $\xi$ . If the first term is  $a_2$ , then the  $n$ -th term is simply  $a_n = n^2$ , since  $a_2$  must be 4. If the first term is  $a_1$ , then the  $n$ -th term is  $a_n = (n + 1)^2$ , since  $ka_1$  must be 4.

The following are all correct representations of the above sequence:

- $\{n^2\}_{n=2}^{\infty}$
- $\{(n + 1)^2\}_{n=1}^{\infty}$
- $\{(n + 2)^2\}_{n=0}^{\infty}$

(c) **The Geometric (Exponential) Pattern.**

If the terms are obtained by *multiplying* by the same number each time, then the  $n$ -th term will look like that number to the  $(n \pm \xi)$ -th power, where the  $\xi$  depends on where the sequence starts.

**Example.**  $\{\frac{1}{2}, 1, 2, 4, 8, 16, \dots\}$ .

Notice that each term in the sequence is twice the one before. Therefore the  $n$ -th term is  $2^{n \pm \xi}$  for some  $\xi$ . If the first term is  $a_{-1}$ , then the  $n$ -th term is simply  $a_n = 2^n$ , since  $a_{-1}$  must be  $\frac{1}{2}$ . If the first term is  $a_1$ , then the  $n$ -th term is  $a_n = 2^{n-2}$ , since  $a_1$  must be  $\frac{1}{2}$ .

The following are all correct representations of the above sequence:

- $\{2^{n-2}\}_{n=1}^{\infty}$
- $\{2^{n+1}\}_{n=-2}^{\infty}$
- $\{2^n\}_{n=-1}^{\infty}$

(d) **The Alternating Pattern.**

If the terms *alternate*, *i.e.* switch from positive to negative every term, then the pattern contains a multiple of  $(-1)^{n \pm \xi}$  for some  $\xi$ . If  $n \pm \xi$  is even, then the  $n$ -th term will be positive, so adjust  $\xi$  so that the correct terms are positive.

**Example.**  $\{\frac{2}{3}, -\frac{2}{3}, \frac{2}{3}, -\frac{2}{3}, \dots\}$ .

The following are all correct representations of the above sequence:

- $\{(-1)^{n-1} \frac{2}{3}\}_{n=1}^{\infty}$
- $\{(-1)^{n+1} \frac{2}{3}\}_{n=1}^{\infty}$
- $\{(-1)^n \frac{2}{3}\}_{n=0}^{\infty}$

(e) **A Combination of Patterns.**

Combine the above techniques, but be careful to adjust everything so that the patterns all start correctly.

**Example.**  $\{\frac{1}{3}, -\frac{5}{9}, \frac{9}{27}, -\frac{13}{81}, \dots\}$ .

Observe the following patterns:

- (i) Counting by 4's in the numerators
- (ii) Powers of 3 in the denominators
- (iii) Alternating terms, starting with a negative

Therefore our  $n$ -th term will look something like

$$(-1)^{\clubsuit} \frac{4n \pm \heartsuit}{3^{\spadesuit}}.$$

Choose an  $n$  to begin with. I'll pick  $n = 0$ . Then adjust everything so that plugging in  $n = 0$  gives  $a_0 = \frac{1}{3}$ :

$$a_n = (-1)^{n+1} \frac{4n+1}{3^n}.$$

Alternate solution: start with  $n = 1$ . Then the sequence is

$$\left\{ (-1)^n \frac{4n-3}{3^{n-1}} \right\}_{n=1}^{\infty}$$

Notice the following convenient trick:

**Convenient Trick.** To start the sequence from  $n = 1$  instead of  $n = 0$ , I replaced all the  $n$ 's in the  $n$ -th term formula by  $n - 1$ . Similarly, if I had wanted to begin with  $n = -10$ , I could have replaced all the  $n$ 's in the  $n$ -th term formula by  $n + 10$ :

$$\left\{ (-1)^{n+11} \frac{4n+41}{3^{n+10}} \right\}_{n=-10}^{\infty}$$

---

**Practice Problems.**

For each problem, find a formula for  $a_n$ . If your first term is not  $a_1$ , be sure to make it clear what your first term is by writing the sequence in the form  $\{a_n\}_{n=?}^{\infty}$ .

1.  $\{1 \cdot 4, 4 \cdot 8, 7 \cdot 16, 10 \cdot 32, \dots\}$

2.  $\{-8, -1, 0, 1, 8, 27, 64, \dots\}$

3.  $\{-\frac{1}{6}, \frac{4}{7}, -2, \frac{64}{9}, -\frac{256}{10}, \dots\}$