

**Section 3.6, Problem 1(b,d):**

Find the orders of each of these permutations.

(b)  $(1, 2, 5)(2, 3, 4)(5, 6)$

(d)  $(1, 2, 3)(2, 4, 3, 5)(1, 3, 2)$

**Solution:**



We know that if a permutation is written as a product of disjoint cycles, then its order is the LCM of the lengths of the cycles. So first we write each permutation as a product of disjoint cycles.

(b)  $(1, 2, 5)(2, 3, 4)(5, 6) = (1, 2, 3, 4, 5, 6)$ . The order of this permutation is 6.

(d)  $(1, 2, 3)(2, 4, 3, 5)(1, 3, 2) = (1, 5, 3, 4)(2) = (1, 5, 3, 4)$ . The order of this permutation is 4.

**Section 3.6, Problem 4:**

Find the permutations that correspond to the rigid motions of a rectangle that is not a square. Do the same for the rigid motions of a rhombus (diamond) that is not a square.

**Solution:**


Rigid motions of the rectangle:  $(1)$ ,  $(1, 4)(2, 3)$ ,  $(1, 2)(3, 4)$ ,  $(1, 3)(2, 4)$ .

Rigid motions of the rhombus:  $(1)$ ,  $(1, 3)$ ,  $(2, 4)$ ,  $(1, 3)(2, 4)$ .

**Section 3.6, Problem 9:**

A rigid motion of a cube can be thought of either as a permutation of its eight vertices or as a permutation of its six sides. Find a rigid motion of the cube that has order 3, and express the permutation that represents it in both ways, as a permutation on eight elements and as a permutation on six elements.

**Solution:**



Rigid motion of order 3: rotation about the line shown (passing through vertices 1 and 7), through an angle of 120 degrees.

Permutation of vertices:  $(2, 4, 5)(3, 8, 6)$ .

Permutation of sides:  $(F, L, B)(T, K, R)$  (letters F, L, B, T, K, and R stand for front, left, bottom, top, back, and right respectively).

### Section 3.6, Problem 13:

List the elements of  $A_4$ .

#### Solution:

Recall that  $A_4$  consists of all even permutations in  $S_4$ .

Elements of  $A_4$  are:  $(1)$ ,  $(1, 2, 3)$ ,  $(1, 3, 2)$ ,  $(1, 2, 4)$ ,  $(1, 4, 2)$ ,  $(1, 3, 4)$ ,  $(1, 4, 3)$ ,  $(2, 3, 4)$ ,  $(2, 4, 3)$ ,  $(1, 2)(3, 4)$ ,  $(1, 3)(2, 4)$ ,  $(1, 4)(2, 3)$ .

(Just checking: the order of a subgroup must divide the order of the group. We have listed 12 elements,  $|S_4| = 24$ , and  $12 \mid 24$ .)

### Section 3.6, Problem 25:

Show that  $S_n$  is isomorphic to a subgroup of  $A_{n+2}$ .

#### Solution:

Hint: define  $\phi : S_n \rightarrow A_{n+2}$  by

$$\phi(\sigma) = \begin{cases} \sigma & \text{if } \sigma \text{ is even,} \\ (\sigma)(n+1, n+2) & \text{if } \sigma \text{ is odd} \end{cases}$$

Show that  $\phi$  is a homomorphism.

### Section 3.7, Problem 3(b):

Show that the following functions are homomorphisms.

(b)  $\phi : \mathbb{R}^\times \rightarrow \mathbb{R}^\times$  defined by  $\phi(x) = \frac{x}{|x|}$ .

#### Solution:

$$\phi(xy) = \frac{xy}{|xy|} = \frac{xy}{|x| \cdot |y|} = \frac{x}{|x|} \frac{y}{|y|} = \phi(x)\phi(y).$$

### Section 3.7, Problem 4:

Let  $G$  be an abelian group, and let  $n$  be any positive integer. Show that the function  $\phi : G \rightarrow G$  defined by  $\phi(x) = x^n$  is a homomorphism.

**Solution:**

Since the group is abelian, any two elements  $x$  and  $y$  commute. Thus  $\phi(xy) = (xy)^n = (xy)(xy) \dots (xy) = xx \dots xyy \dots y = x^n y^n$ .

### Section 3.7, Problem 7(b,d):

Which of the following functions are homomorphisms?

(b)  $\phi : \mathbb{R} \rightarrow \text{GL}_2(\mathbb{R})$  defined by  $\phi(a) = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix}$

(d)  $\phi : \text{GL}_2(\mathbb{R}) \rightarrow \mathbb{R}^\times$  defined by  $\phi\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}\right) = ab$

**Solution:**

(b)  $\phi(a+b) = \begin{bmatrix} 1 & a+b \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & b \\ 0 & 1 \end{bmatrix} = \phi(a)\phi(b)$ , so  $\phi$  is a homomorphism.

(d) Since  $\phi(I_2) = 0 \neq 1$ , the function is not a homomorphism (recall that any homomorphism must send the identity element to the identity element).

### Section 3.7, Problem 9:

Let  $\phi$  be a group homomorphism of  $G_1$  onto  $G_2$ . Prove that if  $G_1$  is abelian then so is  $G_2$ ; prove that if  $G_1$  is cyclic then so is  $G_2$ . In each case, give a counterexample to the converse of the statement.

**Solution:**

Let  $G_1$  be abelian. Let  $a, b \in G_2$ . Since  $\phi$  is onto, there exist  $x, y \in G_1$  such that  $\phi(x) = a$  and  $\phi(y) = b$ . Then  $ab = \phi(x)\phi(y) = \phi(xy) = \phi(yx) = \phi(y)\phi(x) = ba$ , so  $G_2$  is abelian.

Let  $G_1$  be cyclic. Then  $G_1 = \langle x \rangle$  for some  $x \in G_1$ , i.e.  $G_1 = \{x^n \mid n \in \mathbb{Z}\}$ . Then  $G_2 = \phi(G_1) = \{\phi(x^n) \mid n \in \mathbb{Z}\} = \{(\phi(x))^n \mid n \in \mathbb{Z}\} = \langle \phi(x) \rangle$ , so  $G_2$  is cyclic.

Now consider  $\phi : \text{GL}_2(\mathbb{R}) \rightarrow \mathbb{R}^\times$  defined by  $\phi(A) = \det(A)$ . Since  $\text{GL}_2(\mathbb{R})$  is not abelian but  $\mathbb{R}^\times$  is abelian, we have a counterexample to the converse of the first statement.

Finally, consider  $\phi : \mathbb{Z}_2 \times \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$  defined by  $\phi((x, y)) = (x, 0)$ . Since  $\mathbb{Z}_2$  is cyclic, but  $\mathbb{Z}_2 \times \mathbb{Z}_2$  is not, we have a counterexample to the converse of the second statement.

Alternatively,  $\phi : S_3 \rightarrow \mathbb{Z}_2$  defined by

$$\phi(\sigma) = \begin{cases} 0 & \text{if } \sigma \text{ is even,} \\ 1 & \text{if } \sigma \text{ is odd} \end{cases}$$

is a counterexample to the converses of both statements, since  $\mathbb{Z}_2$  is both abelian and cyclic, but  $S_3$  is neither.

(Note: there are many other counterexamples.)