

Practice test 1 - Solutions

Multiple choice questions: circle the correct answer

1. Find the exact value of $\arcsin(1)$.

- A. 0 **(B)** $\frac{\pi}{2}$ C. π D. $\frac{3\pi}{2}$ E. 2π

2. Find the exact value of $\arccos\left(\frac{1}{2}\right)$.

- A. 0 B. $\frac{\pi}{6}$ C. $\frac{\pi}{4}$ **(D)** $\frac{\pi}{3}$ E. $\frac{\pi}{2}$

3. Find the exact value of $\sin\left(\arctan\left(\frac{3}{4}\right)\right)$.

- A. $-\frac{3}{5}$ B. $-\frac{3}{4}$ **(C)** $\frac{3}{5}$ D. $\frac{3}{4}$ E. $\frac{4}{5}$

4. Suppose 100 dollars are invested at an annual interest rate of 10% while interest is compounded monthly. What is the amount after 10 years?

- A. $100\left(1 + \frac{1}{120}\right)^{10}$ **(B)** $100\left(1 + \frac{1}{120}\right)^{120}$ C. $100\left(1 + \frac{10}{12}\right)^{10}$
 D. $120\left(1 + \frac{10}{12}\right)^{100}$ E. $120\left(1 + \frac{1}{120}\right)^{100}$

5. The graph of any exponential function $f(x) = a^x$ (where $a > 0$, $a \neq 1$) passes through which of the following points:

- A. (0, 0) B. (1, 0) **(C)** (0, 1) D. (1, 1)
 E. none of the above

6. If $m(t) = m_0 e^{kt}$ is the mass remaining from an initial mass m_0 of a radioactive substance after time t , find the half-life of the substance.

- A. $m_0/2$ B. $-t/\ln(2)$ C. $k/2$ **(D)** $-\ln(2)/k$ E. $m_0 \ln(2)/k$

Regular problems: show all your work

7. (a) $3x^2y^3 + 3x^3y^2y' - 3y^3 - 9xy^2y' + 4y' = 0$

$$(3x^3y^2 - 9xy^2 + 4)y' = 3y^3 - 3x^2y^3$$

$$y' = \frac{3y^3 - 3x^2y^3}{3x^3y^2 - 9xy^2 + 4}$$

(b) $2^3 - 3 \cdot 2 + 4 = 6$

(c) $y'(2) = \frac{3 - 3 \cdot 2^2}{3 \cdot 2^3 - 9 \cdot 2 + 4} = -\frac{9}{10}$

8. $\tan y + x \sec^2 y \cdot y' + y + xy' + 3y' = 0$

$$(x \sec^2 y + x + 3)y' = -\tan y - y$$


If $x = 0$ and $y = 0$, then $3y'(0) = 0$, so the slope of the tangent line is 0.

9. (a) Let x be the distance between the boy and the point P , let y be the distance between the girl and P , and let z be the distance between the boy and the girl. Then $x^2 + y^2 = z^2$ where x , y , and z are functions of time.

Differentiating this equation with respect to t gives

$$2xx' + 2yy' = 2zz'$$

$$xx' + yy' = zz'$$



45 minutes after the girl started walking (and thus 50 minutes after the boy started walking), $x = 6 \cdot \frac{50}{60} = 5$, $y = 15 - 4 \cdot 4560 = 15 - 3 = 12$, and $z = \sqrt{5^2 + 12^2} = 13$. x' is the rate of change of x , i.e. the speed of the boy, so $x' = 6$, and y' is the rate of change of y , i.e. negative the speed of the girl since y is decreasing, so $y' = -4$. Therefore

$$5 \cdot 6 + 12 \cdot (-4) = 13z'$$

Answer: $-\frac{18}{13}$, decreasing.


- (b) Let x be the distance between the boy and the point P , let y be the distance between the girl and her starting point Q , and let z be the distance between the boy and the girl.

Then $(x + y)^2 + 15^2 = z^2$ (see the figure)

Differentiating this equation with respect to t gives

$$2(x + y)(x' + y') = 2zz'$$

$$(x + y)(x' + y') = zz'$$



45 minutes after the girl started walking (and thus 50 minutes after the boy started walking), $x = 6 \cdot \frac{50}{60} = 5$, $y = 4 \cdot 4560 = 3$, so $x + y = 8$, and $z = \sqrt{8^2 + 15^2} = 17$. x' is the rate of change of x , i.e. the speed of the boy, so $x' = 6$, and y' is the rate of change of y , i.e. the speed of the girl, so $y' = 4$. Therefore

$$(5 + 3)(6 + 4) = 17z'$$

Answer: $\frac{80}{17}$, increasing.

10. $V(t) = \frac{4}{3}\pi(r(t))^3$

$$V'(t) = 4\pi(r(t))^2 r'(t)$$

If $r' = -1$ and $r = 3$, $V'(t) = 4\pi 3^2 \cdot 1 = 36\pi$

Answer: $36\pi \text{ cm}^3/\text{min}$.

11. Since initially there are 800 bacteria, $P(t) = 800e^{kt}$. At $t = 3$ we have: $2700 = 800e^{k \cdot 3}$
 $(e^k)^3 = \frac{27}{8} e^k = \frac{3}{2}$. Then at $t = 5$: $P(5) = 800e^{k \cdot 5} = 800(e^k)^5 = 800 \left(\frac{3}{2}\right)^5 = \frac{800 \cdot 3^5}{2^5} = 25 \cdot 343 = 6075$.

12. (a) $f'(x) = \frac{3}{\sqrt{1-9x^2}}$

(b) $g'(x) = \tan^{-1}(1-x) + \frac{-x}{1+(1-x)^2}$

(c) $h(x) = \frac{-\frac{\sqrt{1-x^2}}{\sqrt{1-x^2}} - \arccos(x) \frac{-2x}{2\sqrt{1-x^2}}}{1-x^2} = \frac{-1 + \frac{x \arccos(x)}{\sqrt{1-x^2}}}{1-x^2}$